

**Quantifying Elephant Social Structure:  
Using a Bilinear Mixed Effects Model to Elicit  
Qualities of Elephant Behavior**

Eric Vance

*Institute of Statistics & Decision Sciences*

**Duke University**

March 7, 2005

## Outline

- Background on elephant social structure
- Data collection and scientific questions
- Peter Hoff's "Bilinear Mixed Effects Models for Dyadic Data" in *JASA* 2005
- Results
- Conclusions
- Future Research

## Elephant Social Structure



- Only females form families. Males just run around looking to mate.
- Oldest female is the leader since she's the largest and wisest.
- Most elephants within a family are related.

## Scientific Questions



- Why do elephants stay in large groups even when food is scarce?
- What role does genetics play in elephant social structure?
- How does one quantify social structure in order to assess whether or not groups are larger in the **Wet Season** vs. the **Dry Season**?

## Data Collection



- Biologists in Kenya ride into the National Park looking for herds of elephants.
- When a herd is spotted, they write down the names of the elephants present.
- The biologists either stay to observe the family or move on to the next herd.

## The Model

- Data is binomial
  - $y_{ij} \sim \text{Bin}(n_{ij}, p_{ij})$
  - $y_{ij}$  is the number of times elephants  $i$  and  $j$  observed together.
  - $n_{ij}$  is the number of times either  $i$  or  $j$  observed.
  
- Use a GLM
  - $E(y_{ij} | \theta_{ij}) = g(\theta_{ij})$ .
  - $g$  is the inverse logit link function.
  - So  $p_{ij} = \frac{\exp \theta_{ij}}{1 + \exp \theta_{ij}}$ .
  
- $\theta_{ij}$  *is the linear predictor.*

## Linear Predictor $\theta_{ij}$

### *How often are elephants together?*

- Intrinsic sociability  $a_i$ .
  - Sociable elephants will be observed together with other elephants (in groups) more often than unsociable elephants.
- Common intercept  $\beta_0$ .
- Genetic relatedness  $\beta_g g_{ij}$ .
  - DNA samples lead to a measure  $g_{ij}$  of how closely elephant  $i$  and  $j$  are related.
- Normal error  $\gamma_{ij}$ .
- **Pairwise effect**  $z_i'z_j$  between elephants  $i$  and  $j$ .


$$\theta_{ij} = \left(\frac{1}{2}\beta_0 + \mathbf{a}_i\right) + \left(\frac{1}{2}\beta_0 + \mathbf{a}_j\right) + \beta_g \mathbf{g}_{ij} + \gamma_{ij} + \mathbf{z}_i' \mathbf{z}_j$$

## Pairwise Effect

$\mathbf{z}_i^T \mathbf{z}_j$  is the inner product of the positions of elephants  $i$  and  $j$  in “Social Space”.

- I fix the dimension of social space  $\mathbf{k} = 2$ .

Elephants  $i$  and  $j$  have positions  $\mathbf{z}_i$  and  $\mathbf{z}_j$  in 2D social space.



$$\mathbf{z}_i \sim N(\mathbf{0}, \sigma_z^2 \mathbf{I})$$

$$\mathbf{z}_j \sim N(\mathbf{0}, \sigma_z^2 \mathbf{I})$$

- If  $\mathbf{z}_i^T \mathbf{z}_j = 0$  then elephants  $i$  and  $j$  interact as often as their sociabilities  $a_i$  and  $a_j$  and their genetics  $g_{ij}$  would predict.
- If  $\mathbf{z}_i^T \mathbf{z}_j > 0$  then  $i$  and  $j$  like each other and are observed together more often than the model would otherwise predict.
- If  $\mathbf{z}_i^T \mathbf{z}_j < 0$  then  $i$  and  $j$  dislike each other.

## Empirical Bayes Priors

$$\theta_{ij} = \left(\frac{1}{2}\beta_0 + \mathbf{a}_i\right) + \left(\frac{1}{2}\beta_0 + \mathbf{a}_j\right) + \beta_g \mathbf{g}_{ij} + \gamma_{ij} + \mathbf{z}'_i \mathbf{z}_j$$

Intercept:  $\beta_0 \sim N(0, \hat{\sigma}_{\beta_0}^2)$

Sociabilities:  $a_i, a_j \sim N(0, \sigma_{\text{soc}}^2), \quad \sigma_{\text{soc}}^2 \sim IG(, )$

Genetic Coefficient:  $\beta_g \sim N(0, \hat{\sigma}_{\beta_g}^2)$

Pairwise error:  $\gamma_{ij} \sim N(0, \sigma_{\gamma}^2), \quad \sigma_{\gamma}^2 \sim IG(, )$

Social space:  $\mathbf{z}_i, \mathbf{z}_j \sim N(\mathbf{0}, \begin{bmatrix} \sigma_z^2 & 0 \\ 0 & \sigma_z^2 \end{bmatrix}), \quad \sigma_z^2 \sim IG(, )$

## Parameter Estimation

***Proceed using Gibbs sampling.***

For details see Peter Hoff's article "Bilinear Mixed-Effects Models for Dyadic Data" in the current March 2005 issue of *JASA*.

## Results


- One elephant Amy
  - Sociability  $a_{Amy}$
  - Position  $\mathbf{z}_{Amy}$  in social space
- Family AA (10 elephants)
  - Is the intercept  $\beta_0$  different in the **Dry** vs. the **Wet** season?
  - What are the elephants' sociabilities?
  - Is the genetic coefficient  $\beta_g$  different from 0 (**Dry** vs. **Wet**)?
  - What does Family AA social space look like?
    - Three elephants Amy, Amelia, and Alison.
    - The **Dry Season** only.
    - All four categories **Dry**, **Wet**, **Dry with Genetics**, and **Wet with Genetics**.

**Amy, Matriarch of Family AA**




# Amy Sociability MCMC


Dry Season




Dry Season with Genetics




Wet Season




Wet Season with Genetics



# Amy Sociability Posterior Density



# Amy Social Space Posterior Draws



## Elephant Family Results




## Revisiting the Model

### *How often are elephants together?*

- Individual sociability  $a_i$ .
  - Sociable elephants will be observed together with other elephants (in groups) more often than unsociable elephants.
- Common intercept  $\beta_0$ .
- Genetic relatedness  $\beta_g g_{ij}$ .
  - DNA samples lead to a measure  $g_{ij}$  of how closely elephant  $i$  and  $j$  are related.
- Normal error  $\gamma_{ij}$ .
- **Pairwise effect**  $z'_i z_j$  between elephants  $i$  and  $j$ .


$$\theta_{ij} = \left(\frac{1}{2}\beta_0 + \mathbf{a}_i\right) + \left(\frac{1}{2}\beta_0 + \mathbf{a}_j\right) + \beta_g \mathbf{g}_{ij} + \gamma_{ij} + \mathbf{z}'_i \mathbf{z}_j$$

# Posterior Intercepts




# Posterior Sociabilities


Dry Season




Dry Season with Genetics




Wet Season




Wet Season with Genetics




# Posteriors for Genetic Coefficients



# Three Elephants' Social Space Posterior Draws



# Dry Season Social Space Posteriors





## Conclusions

- As the matriarch of the family much of Amy's sociability is due to her genetics.
- Some of the posterior sociabilities in Family AA change depending on the season.
- Posterior intercepts  $\beta_0$  for the **Wet** seasons are greater than in the **Dry** seasons, indicating that the elephants are more gregarious during the **Wet** season.
- Genetic coefficients  $\beta_g > 0$  for both **Wet** and **Dry** seasons.
- There are clusters of elephants in social space.

## Future Research

- Include a seasonal intercept (indicator variable) in order to run **Wet** and **Dry** season data sets together.
- Use the real, (partially missing) observation matrices instead of the binomial summarized data.

$$\begin{bmatrix} 1 & 1 & 1 & 1 & \vdots & 0 & 0 \\ 1 & 1 & 1 & 1 & \vdots & 0 & 0 \\ 1 & 1 & 1 & 1 & \vdots & 0 & 0 \\ 1 & 1 & 1 & 1 & \vdots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & \vdots & ? & ? \\ 0 & 0 & 0 & 0 & \vdots & ? & ? \end{bmatrix}$$

## Apply to Lion Social Structure?



## Genetic Relatedness

AA	Amy	Angelina	Amber	Audrey	Alison	Astrid	Agatha	Althea	Amelia
Amy									
Angelina	0.39								
Amber	0.46	0.26							
Audrey	0.31	0.22	0.11						
Alison	0.3	0.02	0.08	0.02					
Astrid	0.25	0.18	0.02	0.14	0.53				
Agatha	0.27	0.08	0.25	-0.15	0.38	0.34			
Althea	0.05	0.05	0.12	-0.01	0.15	0.21	0.46		
Amelia	0.35	0.06	-0.09	0.01	0.09	0.2	0.1	0.02	
Anghared	0.25	0.34	0.02	0.2	-0.06	0.1	0.06	-0.04	0.52

## Family AA Observations

DRY	Amy	Amy	Ang	Ang	Amb	Amb	Aud	Aud	Ali	Ali	Ast	Ast	Aga	Aga	Alt	Alt	Ame	Ame
	P	A	P	A	P	A	P	A	P	A	P	A	P	A	P	A	P	A
Amy P	272	0	237	7	245	2	245	8	154	64	147	61	215	54	205	54	182	68
Amy A	0	159	35	152	27	157	27	151	118	95	125	98	57	105	67	105	90	91
Ang P	237	7	244	0	224	23	217	36	154	64	142	66	189	80	179	80	162	88
Ang A	35	152	0	187	20	164	27	151	90	123	102	121	55	107	65	107	82	99
Amb P	245	2	224	23	247	0	220	33	153	65	141	67	201	68	194	65	174	76
Amb A	27	157	20	164	0	184	27	151	94	119	106	117	46	116	53	119	73	108
Aud P	245	8	217	36	220	33	253	0	149	69	145	63	205	64	197	62	170	80
Aud A	27	151	27	151	27	151	0	178	104	109	108	115	48	114	56	116	83	98
Ali P	154	64	154	64	153	65	149	69	218	0	191	17	147	122	144	115	115	135
Ali A	118	95	90	123	94	119	104	109	0	213	27	196	71	91	74	98	103	78
Ast P	147	61	142	66	141	67	145	63	191	17	208	0	148	121	145	114	121	129
Ast A	125	98	102	121	106	117	108	115	27	196	0	223	60	102	63	109	87	94
Aga P	215	54	189	80	201	68	205	64	147	122	148	121	269	0	257	2	193	57
Aga A	57	105	55	107	46	116	48	114	71	91	60	102	0	162	12	160	76	105
Alt P	205	54	179	80	194	65	197	62	144	115	145	114	257	2	259	0	185	65
Alt A	67	105	65	107	53	119	56	116	74	98	63	109	12	160	0	172	74	107
Ame P	182	68	162	88	174	76	170	80	115	135	121	129	193	57	185	65	250	0
Ame A	90	91	82	99	73	108	83	98	103	78	87	94	76	105	74	107	0	181

